	(FILE	E 'HOME' ENTERED AT 13:08:37 ON 27 FEB 2003)					
L1	FILE 'INPADOC' ENTERED AT 13:08:52 ON 27 FEB 2003 1 S W09519108/PN						
	FILE 2003	'EUROPATFULL, PCTFULL, USPAT2, WPIDS' ENTERED AT 13:30:31 ON 27 FEB					
L2		1651 S CYCLODEXTRIN#(S)OIL#					
L3		156 S L2(S)(SKIN OR HAIR OR TOPICAL OR DERM?)					
	FILE	'USPATFULL' ENTERED AT 13:36:37 ON 27 FEB 2003					
L4		92 S L3					
L5		16 S L4 NOT PY>=1999					
L6		4 S L4/CLM					

BEST AVAILABLE COPY

L5 ANSWER 2 OF 16 USPATFULL

ACCESSION NUMBER: 1998:138878 USPATFULL

TITLE: Process for preparing decolorized carotenoid-

cyclodextrin complexes

INVENTOR(S): Sikorski, Christopher, 1805 Davis St., Whiting, IN,

United States 46394

Schwartz, Joel L., 5225 Pooks Hill, Promenade Apt. 315

South, Bethesda, MD, United States 20814

Shklar, Gerald, 7 Chauncy La., Cambridge, MA, United

States 02138

NUMBER KIND DATE

PATENT INFORMATION: APPLICATION INFO.:

US 5834445 19981110 US 1995-552374 19951103 (8)

RELATED APPLN. INFO.: Continuation-in-part of Ser. No. US 1994-339018, filed

on 14 Nov 1994, now abandoned which is a continuation-in-part of Ser. No. US 1992-947067, filed

on 18 Sep 1992, now abandoned which is a continuation of Ser. No. US 1991-741203, filed on 30 Jul 1991, now abandoned which is a continuation of Ser. No. US

1990-469171, filed on 24 Jan 1990, now abandoned And a continuation-in-part of Ser. No. US 1989-392857, filed

on 11 Aug 1989, now abandoned , said Ser. No. US 1992-947067, filed on 18 Sep 1992, now abandoned which

BEST AVAILABLE COPY

is a continuation-in-part of Ser. No. US 1992-860201, filed on 26 Mar 1992, now abandoned which is a

continuation of Ser. No. US 1991-708130, filed on 29

May 1991, now abandoned

DOCUMENT TYPE: FILE SEGMENT: Utility Granted

PRIMARY EXAMINER: LEGAL REPRESENTATIVE: Henley, III, Raymond Trask, Britt & Rossa

NUMBER OF CLAIMS: EXEMPLARY CLAIM:

18 1

2 Drawing Figure(s); 2 Drawing Page(s)

NUMBER OF DRAWINGS: 2 Dr LINE COUNT: 1150

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

DETD . . . to 50 milligrams of vitamin E (as DL-alpha tocopherol acid succinate). To this mixture were added 50 milligrams of .beta.

cyclodextrin and 50 milligrams of gelatin derived from bovine skin (bloom 225, type III). The color of the mixture was dark brown-red, the texture was coarse and sandy. Next, 1. . . the

texture

the

of the mixture to a somewhat watery-creamy texture, but the color remained unchanged. The composition so prepared felt **oily** on the **skin** and contained small but visible particles of Vitamin E, and .beta. carotene particles suspended in solution.

DETD Fifty milligrams of .beta. carotene were added to 100 milligrams of .beta. cyclodextrin and 1 milliliter of d-H.sub.2 O. The

.beta. carotene was in a suspension, the color of the mixture was light.

. . color and did not change the texture. Some suspended particles of Vitamin E were observed. One milliliter of light mineral  ${\tt oil}$  further increased the solubilization of both the .beta. carotene and

vitamin E. The color of the composition was yellow-tan. . . watery. Glycerine (1 milliliter) and/or gelatin (50 mg) were added to the mixture, followed by 1 milliliter of LUBRIDERM brand skin

cream. (LUBRIDERM has the following ingredients in order of amount: water, mineral oil, petrolatum, sorbitol, lanolin, lanolin alcohol, triethanolamine, cetyl alcohol, butylparaben, methylparaben, propylparaben, sodium chloride.) The result was a cream with a. DETD . examples it is readily seen that it is highly preferable to solubilize the .beta. carotene by combining it with the cyclodextrin, prior to mixing with the other ingredients such as glycerine, mineral oil, or skin cream base. It is further preferable to solubilize the vitamin E succinate prior to mixing with other ingredients. DETD TABLE V Comparative solubility in skin cream of .beta. carotene with different solubilization mixtures Solvent #1 Solvent #2 .beta.CD.sup.1 Gelatin Mineral Oil Glycerin H.sub.2 O H.sub.2 0 7-8 2 3 3 1 H.sub.2 O + DMSO.sup.2

4

Solvent #2. This mixture was in turn combined with LUBRIDERM skin

2

carotene was mixed with Solvent #1 prior to mixing

4

3

cream, and the results scored.

8

H.sub.2 O + DMSO. . .

with

1 = .beta. cyclodextrin

2 = Dimethylsulfoxide

3 = 1,2 Dimethoxyethane

DEST AVAILABLE COPY

```
=> s wo9519108/pn
L1 1 WO9519108/PN
```

=> d fam l1

## L1 ANSWER 1 OF 1 INPADOC COPYRIGHT 2003 EPO

## PATENT FAMILY INFORMATION AN 40905863 INPADOC

+PRAI-		+	+AI	+
US 1991-805724	B2			
US 1993-40052	Α	19930330	US 1993-40052 A 199303	330
			WO 1994-US401 A 199401	12
US 1993-40052	A2	19930330	US 1994-212856 A 199403	315
US 1993-76276	A	19930702	WO 1994-US401 A 199401	12
US 1993-76276	A2	19930702	US 1994-212856 A 199403	315
US 1993-805724	A	19930917	WO 1994-US401 A 199401	12
US 1993-805724	B2	19930917	US 1994-212856 A 199403	315
US 1994-212856	Α	19940315	US 1994-212856 A 199403	315
			WO 1994-US401 A 199401	12
WO 1994-US401	Α	19940112	WO 1994-US401 A 199401	12
WO 1994-US401	W	19940112	EP 1994-906601 A 199401	12
			JP 1994-518992 A 199401	12
			US 1994-212856 A 199403	15
			+PI	
EP 1994-906601	Α	19940112		-
			EP 785715 A4 199707	130
	Α	19940112	JP 09511231 T2 199711	11
US 1993-40052	Α	19930330	US 5296472 A 199403	122
US 1994-212856	A	19940315	US 5380711 A 199501	10
WO 1994-US401	Α	19940112	WO 9519108 A1 199507	120

10 priorities, 5 applications, 6 publications

**BEST AVAILABLE COPY**